

SN 10/656,358
ATTY DOCKET NO. S-100,543

AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:

LISTING OF CLAIMS:

1. (Currently Amended) A method of fluorescence-based cycle sequencing of a sample DNA, comprising,

(a) preparing a reaction mixture containing:

- (i) the sample DNA,
- (ii) a primer set complementary to DNA primer sites flanking or interspersed within the sample DNA, wherein the T_d of the primers in the primer set are between about 72 °C and 75 °C,
- (iii) a thermostable polymerase,
- (iv) a mixture of dNTPs and fluorescently-labeled ddNTPs, and
- (v) a suitable buffer

(b), dissociating the sample DNA to create single stranded templates, wherein said dissociation is achieved by heating the sample DNA to between about 92 °C and 95 °C for ~~at least~~ about 3 minutes;

(c) annealing the primers to the primer sites, wherein said annealing is achieved at a temperature of between about 65°C and 67°C for ~~at least~~ about 30 seconds;

(d) extending the annealed primers to generate a series of fluorescently-labeled dideoxynucleic acid fragments, wherein said primer extension is achieved at a temperature of between about 75°C and 78°C for between about 3 to 4 minutes;

SN 10/656,358
ATTY DOCKET NO. S-100,543

(e) heating the reaction mixture to between about 92°C and 95°C in order to dissociate double stranded DNA;

(f) repeating the steps c through e for a plurality of cycles; and

(eg) determining the nucleotide sequence of the sample DNA from the series of fluorescently-labeled dideoxynucleic acid fragments present in the reaction mixture.

2. (Currently Amended) The method according to claim 1, wherein the number of cycles is between about 30 and 50 cycles.
3. (Currently Amended) The method according to claim 1, wherein the number of cycles is between about 50 and 60 cycles.
4. (Currently Amended) The method according to claim 1, wherein the number of cycles is between about 60 and 70 cycles.
5. (Currently Amended) The method according to claim 1, wherein the ~~the~~ primers are complementary to a PUC18 vector containing the sample DNA and have the following nucleotide sequences:

5' GCT GCA AGG CGA TTA AGT TGG GTA 3' (SEQ ID NO: 1)

5' GTT GTG TGG AAT TGT GAG CGG ATA AC 3' (SEQ ID NO: 2)

SN 10/656,358
ATTY DOCKET NO. S-100,543

6-10. (Original)

THIS PAGE BLANK (USPTO)